

Routine childhood immunisations from Autumn 2006

When to immunise	Diseases protected against	Vaccine given	Immunisation site
Two months old	Diphtheria, tetanus, pertussis, polio and <i>Haemophilus influenzae</i> type b (Hib)	DTaP/IPV/Hib (PediaceL)	Thigh
	Pneumococcal infection	PCV (Prevenar)	Thigh
Three months old	Diphtheria, tetanus, pertussis, polio and <i>Haemophilus influenzae</i> type b (Hib)	DTaP/IPV/Hib (PediaceL)	Thigh
	Meningitis C	MenC (Menjugate, Meningitec or Neisvac C)	Thigh
Four months old	Diphtheria, tetanus, pertussis, polio and <i>Haemophilus influenzae</i> type b (Hib)	DTaP/IPV/Hib (PediaceL)	Thigh
	Meningitis C	MenC (Menjugate, Meningitec or Neisvac C)	Thigh (2.5cm from PediaceL injection)
	Pneumococcal infection	PCV (Prevenar)	Opposite thigh
Around 12 months old	Hib/MenC	Hib/MenC (Menitorix)	Thigh
Around 13 months old	Measles, mumps and rubella	MMR (Priorix or MMR II)	Thigh
	Pneumococcal infection	PCV (Prevenar)	Thigh
Three years four months to five years old	Diphtheria, tetanus, pertussis and polio (dTaP/IPV or DTaP/IPV)	dTaP/IPV (Repevax) or DTaP/IPV (Infanrix-IPV)	Upper arm
	Measles, mumps and rubella	MMR (Priorix or MMR II) (check first dose has been given)	Upper arm
Thirteen to 18 years old	Tetanus, diphtheria and polio	Td/IPV (Revaxis), and check MMR status	Upper arm



All vaccines for use in the childhood immunisation programme are supplied free of charge by Healthcare Logistics on 0870 871 1890

*NB Where a vaccine is manufactured by more than one supplier it may, on occasion, be necessary to substitute an alternative brand.