

GS12 Laparoscopic Inguinal Hernia Repair (TAPP)

What is an inguinal hernia?

An inguinal hernia is a common type of hernia, causing a lump and sometimes pain in the groin.

Your surgeon has recommended a transabdominal preperitoneal (TAPP) hernia operation. However, it is your decision to go ahead with the operation or not. This document will give you information about the benefits and risks to help you make an informed decision.

If you have any questions that this document does not answer, you should ask your surgeon or any member of the healthcare team.

How does a hernia happen?

The abdominal cavity contains the intestines and other structures. These are protected by the abdominal wall, which is made up of four layers. The inner layer is a membrane. The second layer is a wall made of muscle. A layer of fat separates the muscle from the outer layer of skin.

Weak spots can develop in the layer of muscle, resulting in the contents of the abdomen, along with the inner layer, pushing through the abdominal wall. This produces a lump called a hernia (see figure 1).




Figure 1

Hernia – bowel pushing through a weakness in the muscle wall of the abdomen

An inguinal hernia happens at the inguinal canal (see figure 2). This is a narrow passage in which blood vessels pass through the abdominal wall .




Figure 2

Position of a left inguinal hernia

The inguinal canal is prone to hernias, which can be due to a defect from birth or from gradual weakening of the muscles.

What are the benefits of surgery?

You should no longer have the hernia. Surgery should prevent you from having any serious complications that a hernia can cause and allow you to return to normal activities.

Are there any alternatives to surgery?

Surgery is recommended as it is the only dependable way to cure the condition. You can sometimes control the hernia with a truss (padded support belt) or simply leave it alone. It will not go away without an operation.

What will happen if I decide not to have the operation?

Hernias will get bigger with time. They can also be dangerous because the intestines or other structures within the abdomen can get trapped and have their blood supply cut off (strangulated hernia). This is serious and needs an urgent and bigger operation, with a higher risk of serious complications. If left untreated, a strangulated hernia can cause death.

What does the operation involve?

Inguinal hernias can be repaired using the laparoscopic ('keyhole') technique or by an open cut in the groin. Your surgeon has recommended a laparoscopic operation for you, as this is associated with less pain, less scarring and a faster return to normal activities.

The operation is performed under a general anaesthetic and usually takes about half an hour (less than an hour for a repair to both sides).

Your surgeon will make a small cut in or near your umbilicus so they can insert an instrument which inflates the abdominal cavity with gas (carbon dioxide). They will make two small cuts on your abdomen so they can insert tubes (ports) into your abdomen. Your surgeon will place surgical instruments through the ports along with a telescope so they can see inside your abdomen and perform the operation.

Your surgeon will return the part of the abdomen showing out and causing the hernia, and insert a synthetic mesh to cover the weak spot. They will close the small cut and holes with stitches or glue.

In about 1 in 100 people it will not be possible to complete the operation using this technique. If this happens, the operation will be changed (converted) to an open procedure.

What should I do about my medication?

You should continue your normal medication unless you are told otherwise. Let your surgeon know if you are on **warfarin** or **clopidogrel**. Follow your surgeon's advice about stopping this medication before the operation.

What can I do to help make the operation a success?

• Lifestyle changes

If you smoke, try to stop smoking now. Stopping smoking several weeks or more before an operation may reduce your chances of getting complications and will improve your long-term health.

For help and advice on stopping smoking, go to www.gosmokefree.co.uk.

You have a higher chance of developing complications if you are overweight.

For advice on maintaining a healthy weight, go to www.eatwell.gov.uk.

• Exercise

Regular exercise can reduce the risk of heart disease and other medical conditions, improve how your lungs work, boost your immune system, help you to control your weight and improve your mood. Exercise should help to prepare you for the operation, help with your recovery and improve your long-term health.

For information on how exercise can help you, go to www.eidoactive.co.uk.

Before you start exercising, you should ask a member of the healthcare team or your GP for advice.

What complications can happen?

The healthcare team will try to make your operation as safe as possible. However, complications can happen. Some of these can be serious and can even cause death. You should ask your doctor if there is anything you do not understand. Any numbers which relate to risk are from studies of people who have had this operation. Your doctor may be able to tell you if the risk of a complication is higher or lower for you.

The complications fall into three categories.

- 1 Complications of anaesthesia
- 2 General complications of any operation
- 3 Specific complications of this operation

1 Complications of anaesthesia

Your anaesthetist will be able to discuss with you the possible complications of having an anaesthetic.

2 General complications of any operation

- **Pain**, which happens with every operation. The healthcare team will try to reduce your pain. They will give you medication to control the pain and it is important that you take it as you are told so you can move about and cough freely. After a laparoscopy, it is common to have some pain in your shoulders because a small amount of gas may be left under the diaphragm. Your body will usually absorb the gas naturally over the next 24 hours, which will ease the symptoms.

- **Bleeding** during or after surgery. This does not usually need a blood transfusion or another operation but it is common to get bruising of the groin and, **in men**, bruising of the penis or scrotum.

- **Infection in the surgical wound**, which may need treatment with antibiotics.

- **Unsightly scarring** of the skin.

- **Blood clots** in the legs (deep-vein thrombosis), which can occasionally move through the bloodstream to the lungs (pulmonary embolus), making it difficult for you to breathe. Nurses will encourage you to get out of bed soon after surgery and may give you injections to reduce the risk of blood clots.

3 Specific complications of this operation

a Laparoscopic complications

- **Damage to internal organs** when placing instruments into the abdomen (risk: 1 in 1,000). The risk is higher in people who have previously had surgery to the abdomen. If an injury does happen, you may need open surgery, which involves a much bigger cut. About 1 in 3 of these injuries is not obvious until after surgery, so if you have pain which does not continue to improve each day after surgery, you should let your doctor know.

- **Developing a hernia** near one of the cuts used to insert the ports (risk: 2 in 10,000). Your surgeon will try to reduce this risk by using small ports (less than 10 millimetres in diameter) where possible or, if they need to use larger ports, using deeper stitching techniques to close the cuts.

- **Injury to the bowel** (risk: 1 in 200). This can happen during surgery or if the bowel becomes trapped between the mesh and the abdominal wall, causing bowel obstruction. You may need further surgery.

- **Surgical emphysema** (crackling sensation in the skin due to trapped gas), which settles quickly and is not serious.

b Hernia repair complications

- **Developing a lump** at the site of the original hernia (risk: 1 in 10). This is caused by a collection of blood or fluid and normally settles over a few weeks.

- **Discomfort or pain** in the groin.

- **In men, discomfort or pain** in the testicle on the side of the operation.

- **In men, difficulty passing urine**. You may need a catheter (tube) in your bladder for a day or two (risk: 1 in 100).

- **In men, damage to the blood supply of the testicle** (risk: less than 1 in 100). This may result in a small non-functioning testicle on the side of the operation.

How soon will I recover?

• In hospital

After the operation you will be transferred to the recovery area and then to the ward. You should be able to go home the same day or the day after. However, your doctor may recommend that you stay a little longer. If you do go home the same day, **a responsible adult should take you home in a car or taxi, and stay with you for at least 24 hours.**

If you are worried about anything, in hospital or at home, ask a member of the healthcare team. They should be able to reassure you or identify and treat any complications.

• Returning to normal activities

You may return to normal activities when you feel comfortable to do so, usually after one week. You do not need to avoid lifting, but you may find it uncomfortable if you lift heavy weights within the first two to four weeks.

Regular exercise should help you to return to normal activities as soon as possible. Before you start exercising, you should ask a member of the healthcare team or your GP for advice.

Do not drive until you are confident about controlling your vehicle and always check with your doctor and insurance company first.

• The future

Most people make a full recovery and can return to normal activities.

Occasionally the hernia comes back (risk: less than 3 in 100). This can happen many years later and may need another operation.

Summary

An inguinal hernia is a common condition caused by a weakness in the abdominal wall, near the inguinal canal. If left untreated, an inguinal hernia can occasionally cause serious complications. Surgery is usually safe and effective. However, complications can happen. You need to know about them to help you make an informed decision about surgery. Knowing about them will also help to detect and treat any problems early.

Further information

- NHS smoking helpline on 0800 169 0 169 and at www.gosmokefree.co.uk
- www.eatwell.gov.uk – for advice on maintaining a healthy weight
- www.eidoactive.co.uk – for information on how exercise can help you
- www.aboutmyhealth.org - for support and information you can trust
- NHS Direct on 0845 46 47 (0845 606 46 47 - textphone)

Acknowledgements

Author: Mr Ian Beckingham DM FRCS

Illustrations: Hannah Ravenscroft RM and LifeART
image copyright 2008 Lippincott Williams & Wilkins.
All rights reserved.

Local information

You can get information locally from your Consultant.
The Trust's switchboard number is 0844 811 8110.
You can also contact:

.....
.....
.....

Tell us how useful you found this document at www.patientfeedback.org

This document is intended for information purposes only and should not replace advice that your relevant health professional would give you.

GS12

Issued December 2008

Expires end of December 2009



www.rcsed.ac.uk



www.asgbi.org.uk

