

SEPSIS 6 TREATMENT PROTOCOL

Patient name
IW number
DOB

Action	Date	Time	Signed	Reason (if not completed)
Give High Flow Oxygen (Give oxygen if needed to keep SpO ₂ 94–98% and if COPD seek advice)				
Give IV Fluids (If Systolic below 90mmHg give 20ml/Kg 0.9% Saline or Hartmanns solution otherwise give 500ml of same fluid)				
Take Blood Cultures				
Take Blood Gas (for Lactate)				
Give Broad Spectrum IV Antibiotics				
Measure Urine Output (Strict fluid balance and may need catheter)				
Apply sky-blue Sepsis wristband				
Inform critical care outreach service for sepsis liaison on bleep 006				

SEPSIS 6 TREATMENT PROTOCOL

Patient name
IW number
DOB

Action	Date	Time	Signed	Reason (if not completed)
Give High Flow Oxygen (Give oxygen if needed to keep SpO ₂ 94–98% and if COPD seek advice)				
Give IV Fluids (If Systolic below 90mmHg give 20ml/Kg 0.9% Saline or Hartmanns solution otherwise give 500ml of same fluid)				
Take Blood Cultures				
Take Blood Gas (for Lactate)				
Give Broad Spectrum IV Antibiotics				
Measure Urine Output (Strict fluid balance and may need catheter)				
Apply sky-blue Sepsis wristband				
Inform critical care outreach service for sepsis liaison on bleep 006				

SEPSIS 6 TREATMENT PROTOCOL

Patient name
IW number
DOB

Action	Date	Time	Signed	Reason (if not completed)
Give High Flow Oxygen (Give oxygen if needed to keep SpO ₂ 94–98% and if COPD seek advice)				
Give IV Fluids (If Systolic below 90mmHg give 20ml/Kg 0.9% Saline or Hartmanns solution otherwise give 500ml of same fluid)				
Take Blood Cultures				
Take Blood Gas (for Lactate)				
Give Broad Spectrum IV Antibiotics				
Measure Urine Output (Strict fluid balance and may need catheter)				
Apply sky-blue Sepsis wristband				
Inform critical care outreach service for sepsis liaison on bleep 006				

SEPSIS 6 TREATMENT PROTOCOL

Patient name
IW number
DOB

Action	Date	Time	Signed	Reason (if not completed)
Give High Flow Oxygen (Give oxygen if needed to keep SpO ₂ 94–98% and if COPD seek advice)				
Give IV Fluids (If Systolic below 90mmHg give 20ml/Kg 0.9% Saline or Hartmanns solution otherwise give 500ml of same fluid)				
Take Blood Cultures				
Take Blood Gas (for Lactate)				
Give Broad Spectrum IV Antibiotics				
Measure Urine Output (Strict fluid balance and may need catheter)				
Apply sky-blue Sepsis wristband				
Inform critical care outreach service for sepsis liaison on bleep 006				